



INTERNATIONAL JOURNAL OF ADVANCE RESEARCH, IDEAS AND INNOVATIONS IN TECHNOLOGY

ISSN: 2454-132X

Impact Factor: 6.078

(Volume 8, Issue 3 - V8I3-1161)

Available online at: <https://www.ijariit.com>

Numerical Solution to the One-Dimensional Heat Equation by using a Recursive Form of Cubic B-spline Collocation Method

Rajashekhar Reddy

yrsreddy4@gmail.com

Jawaharlal Nehru Technological University, Hyderabad, Telangana

ABSTRACT

In this paper, recursive form of quadratic B-spline Collocation method is proposed for calculating numerical solution of one-dimensional heat equation. Recursive form of B-spline is used for the spatial coordinates whereas forward difference scheme is applied for the time derivative. The performance of the method is tested at different time level. The numerical result shows that the present method is a successful numerical technique to find solution for time dependent problems.

Keywords— One Dimensional Heat Equation, B-spline, Collocation Method, Forward difference scheme

1. INTRODUCTION

Heat transfer problems play the important role in boilers, condensers, air pre-heaters, economizers, electric motors, generators and transformers and in chemical reactions. The study of heat transfer problems gives the solutions to optimize the usage of materials and their performance in different systems [7]. Many numerical methods are developed to solve such type of Heat transfer problems. Widely used such numerical methods are finite volume method and finite element method [1, 2, 3, 4]. These methods depend on discretization of domains, conversion of strong form into weak form of the governing equation and evaluation of integrals to obtain linear system of equations. This process leads to errors like geometrical error, reduction in the continuity requirement of the approximating function which demands the fine mesh for an acceptable solution. In addition to these errors, mesh generation is more time consuming and costly.

Owing to the above-mentioned difficulties in the mesh-based methods, various point wise approximation techniques are developed like Smoothed Particle Hydrodynamics (SPH) method, The Element Free Galerkin (EFG), Collocation methods etc. In collocation method, an approximating function is defined based on the nodal distribution only. In this method, many intermediary evaluations such as conversion of strong form differential equation to weak form, evaluation of integrals can be avoided. The use of B-Spline basis functions in collocation method improves the smoothness and accuracy of the solution. In this manuscript, a methodology is developed for the solution of one-dimensional Heat problems using quadratic B-spline Collocation method. Applicability and convergence of the present method is tested by considering numerical heat transfer problem.

One dimensional heat equation is given as

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}, \quad a \leq x \leq b, \quad t \geq 0 \quad (1)$$

subject to the initial condition

$$u(x,0) = f(x), \quad x \in [a,b] \quad (2)$$

and the boundary conditions

$$u(a,t) = g_0(t), \quad u(b,t) = g_1(t), \quad t \in [0,T] \quad (3)$$

2.B-SPLINE COLLOCATION METHOD

The solution domain $a \leq x \leq b$ is partitioned into a mesh of uniform length $h = x_{j+1} - x_j$, where $j=0,1,2,\dots, N-1,N$. Such that $a = x_0 < x_1 < x_2 \dots < x_{n-1} < x_n = b$.

In the cubic B-spline collocation method the approximate solution is written as the linear combination of cubic B-spline basis functions for the approximation space under consideration. The proposed numerical solution for solving Eq. (1)

using the collocation method with cubic B-spline is to find an approximation solution $U^k(x,t)$ at k^{th} time level to the exact solution $U(x,t)$ in the form:

$$U^k(x,t) = \sum_{i=-3}^{n+3} C_i^k(t) N_{i,p}(x) \tag{4}$$

where $C_i(t)$'s are time dependent quantities to be determined from the initial and boundary conditions and collocation from the differential equation.

A zero degree and other than zero degree B-spline basis functions [5, 6] are defined at x_j recursively over the knot vector space $X = \{x_1, x_2, x_3, \dots, x_{n-1}, x_n\}$ as

i) if $p=0$

$$N_{i,p}(x) = 1 \quad \text{if } x \in (x_i, x_{i+i})$$

$$N_{i,p}(x) = 0 \quad \text{if } x \notin (x_i, x_{i+i})$$

ii) if $p \geq 1$

$$N_{i,p}(x) = \frac{x - x_i}{x_{i+p} - x_i} N_{i,p-1}(x) + \frac{x_{i+p+1} - x}{x_{i+p+1} - x_{i+1}} N_{i+1,p-1}(x) \tag{5}$$

where p is the degree of the B-spline basis function and x is the parameter belongs to X . When evaluating these functions, ratios of the form $0/0$ are defined as zero.

Derivatives of B-splines

If $p=3$, we have

$$N'_{i,p}(x) = \frac{x - x_i}{x_{i+p} - x_i} N'_{i,p-1}(x) + \frac{N_{i,p-1}(x)}{x_{i+p} - x_i} + \frac{x_{i+p+1} - x}{x_{i+p+1} - x_{i+1}} N'_{i+1,p-1}(x) - \frac{N_{i+1,p-1}(x)}{x_{i+p+1} - x_{i+1}}$$

$$N^{iii}_{i,p}(x) = 3 \frac{N^{ii}_{i,p-1}(x)}{x_{i+p} - x_i} - 3 \frac{N^{ii}_{i+1,p-1}(x)}{x_{i+p+1} - x_{i+1}} \tag{6}$$

$$(U^k)^{ii}(x) = \sum_{i=-3}^{n+3} C_i^k N^{ii}_{i,p}(x) \tag{7}$$

3. SOLUTION OF HEAT EQUATION

We discrete the time derivative of Eq (1) by a first order accurate forward difference formula and

$$\frac{u^{k+1} - u^k}{\Delta t} = \frac{1}{2} \left(\frac{\partial^2 u^{k+1}}{\partial x^2} + \frac{\partial^2 u^k}{\partial x^2} \right)$$

$$2 * u^{k+1} - \Delta t * \frac{\partial^2 u^{k+1}}{\partial x^2} = 2 * u^k + \Delta t * \frac{\partial^2 u^k}{\partial x^2} \tag{8}$$

Substituting the Eq (2) in Eq (8) then we have

$$2 * \sum_{i=-3}^{n+3} C_i^{k+1}(t) N_{i,p}(x) - \Delta t * \sum_{i=-3}^{n+3} C_i^{k+1}(t) N''_{i,p}(x) =$$

$$2 * \sum_{i=-3}^{n+3} C_i^k(t) N_{i,p}(x) + \Delta t * \sum_{i=-3}^{n+3} C_i^k(t) N''_{i,p}(x) \tag{9}$$

Evaluating Eq. (9) at the nodal points $x = x_0, x_1, x_2, \dots, x_{N-1}, x_N$. These are also collocation points. (n+1) equations are obtained in (n+3) unknowns. Two more equations are obtained by using the boundary conditions. Finally, (n+3) equations are obtained in (n+3) unknowns C_i^{k+1} 's at (k+1)th time level. These unknowns are determined by using the above obtained (n+3) equations. Substituting all the C_i^{k+1} in equation (4), the solution is known to find values $U(x, t)$ at any time in the given domain of x .

3.1 Initial vector

Initial vector $C_i^{0'}$ is obtained by using the initial condition $u(x, 0) = f(x)$ for $x = x_0, x_1, x_2, \dots, x_{N-1}, x_N$ and by the boundary conditions $u_x(x_0, 0) = f'(x)$, $u_x(x_N, 0) = f'(x)$. This yields the (n+3) equations in (n+3) unknowns.

3.2 Numerical Example

One dimensional heat equation is given with initial and boundary conditions as

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}, \quad 0 < x < 1, \quad t \geq 0 \tag{1}$$

subject to the initial condition

$$u(x, 0) = \sin \pi x \quad x \in [0, 1] \tag{2}$$

and the boundary conditions

$$u(0, t) = 0, \quad u(1, t) = 0 \tag{3}$$

The exact solution is known to be $u(x, t) = e^{-\pi^2 t} \sin \pi x$

Table1: Maximum Absolute Relative Error obtained for the problem

Δt	Present method
.0100	8.1199e-004
.0200	1.5X10 ⁻³
.0300	2X10 ⁻³
.0400	2.4X10 ⁻³




Figure1: Comparison of cubic B-spline collocation solution with exact solution for different time levels

4. CONCLUSION

The effectiveness of the proposed method is presented by considering one numerical example. The second-degree B-spline basis function is used in collocation method and forward difference scheme is applied to discretize the time derivative function. Numerical results got by the present method are well agreed with the analytical solution values. By using this method, amount of work time is reduced majorly. This method may be applied to different types of some more time dependent boundary value problems for its efficient.

5. REFERENCES

[1] K.Hollig, Finite Element Methods with B-Splines, The SIAM series on Frontiers in Applied Mathematics.2003.
 [2] B-spline Finite Elements. Computer Methods in Applied Mechanics and Engineering,100

- [3] 325- 337.[http://dx.doi.org/10.1016/0045-7825\(92\)90088-2](http://dx.doi.org/10.1016/0045-7825(92)90088-2)
- [4] Aziz, T., Khan, A. and Rashidinia, J. (2005) Spline Methods for the Solution of Fourth-Order Parabolic Partial Differential Equations. Applied Mathematics and Computation, 167, 153-166.
<http://dx.doi.org/10.1016/j.amc.2004.06.095>
- [5] Caglar, N. and Caglar, H. (2006) B-Spline Solution of Singular Boundary Value Problems. Applied Mathematics and Computation, 182, 1509-1513. <http://dx.doi.org/10.1016/j.amc.2006.05.035>
- [6] H. B. CURRY AND I. J. SCHOENBERG On Polya frequency functions IV: The fundamental spline functions and their limits, J. Anal. Math. 17 (1966), 71-107.
- [7] CARL DE BOOR On Calculating with B-splines. JOURNAL OF APPROXIMATION THEORY 6, SO-62 (1972)
- [8] R.W. Lewis P. Nithiarasu K.N. Seetharamu, Fundamentals of the Finite Element Method for Heat and Fluid Flow 2004 John Wiley & Sons